



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Targeting toll-like receptor 7/8 improves host anti-infective response in alcoholic cirrhosis

Albano, Emanuele ; Stickel, Felix

DOI: <https://doi.org/10.1136/gutjnl-2017-314437>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-144844>

Journal Article

Accepted Version

Originally published at:

Albano, Emanuele; Stickel, Felix (2018). Targeting toll-like receptor 7/8 improves host anti-infective response in alcoholic cirrhosis. *Gut*, 67(10):1749-1750.

DOI: <https://doi.org/10.1136/gutjnl-2017-314437>

**TARGETING TOLL-LIKE RECEPTOR 7/8 IMPROVES HOST
ANTI-INFECTIVE RESPONSE IN ALCOHOLIC CIRRHOSIS**

Journal:	<i>Gut</i>
Manuscript ID	gutjnl-2017-314437.R1
Article Type:	Commentary
Date Submitted by the Author:	n/a
Complete List of Authors:	ALBANO, EMANUELE; University A. Avogadro of East Piedmont, Dept. of Medical Sciences Stickel, Felix; University of Zürich, Department of Gastroenterology and Hepatology; Klinik Beau-Site, Hirslanden, Hepatology Unit
Keywords:	ALCOHOL

Albano & Stickel

toll-like receptor targeting in alcoholic cirrhosis

COMMENTARY

TARGETING TOLL-LIKE RECEPTOR 7/8 IMPROVES HOST ANTI-INFECTIVE RESPONSE IN ALCOHOLIC CIRRHOSIS

Emanuele Albano ¹ & Felix Stickel ²

¹Department of Health Sciences and Interdisciplinary Research, Centre for Autoimmune Diseases, University of East Piedmont, Novara, Italy

²Department of Gastroenterology and Hepatology, University Hospital of Zurich, Zurich, Switzerland

Word count: 875

References: 12

Short title: Toll-like receptor targeting in alcoholic cirrhosis

Correspondence

Emanuele Albano, Dept. of Health Sciences, University of East Piedmont, Via Solaroli 17, 28100 Novara, Italy. Tel: +390321660642; Fax: +390321620421; E-mail: emanuele.albano@med.uniupo.it

The production of reactive oxygen species (ROS) by activated macrophages and neutrophil granulocytes represents an important cause of alcohol-induced oxidative stress and significantly contributes to the pathogenesis of alcoholic liver disease (ALD) [1]. Furthermore, diffuse neutrophil infiltration of the liver is a key feature of acute alcoholic hepatitis actively participating to parenchymal injury [2]. However, during ALD progression to cirrhosis, neutrophil functions such as ROS production, bacterial phagocytosis and granule exocytosis are impaired leading to an increased susceptibility to bacterial infections of cirrhotic patients [3]. Indeed, the development of bacterial peritonitis is a common complication of cirrhosis, while sepsis is a major cause of mortality of patients with decompensated alcoholic cirrhosis being also associated with multi-organ failure and immune deficiencies [4].

The key role of neutrophils in anti-bacterial defences mainly relies on the capacity of the enzyme NADPH oxidase 2 (NOX2) to generate superoxide anion (O_2^-) during a process known as oxidative burst. Superoxide anion, in fact, by conversion to hydrogen peroxide fuels myeloperoxidase-mediated production of the powerful bactericidal agents, hypochlorite and chloramine [5]. NOX2 is a multi-protein enzymatic system consisting of the transmembrane proteins gp91^{phox} and p22^{phox} forming the catalytic unit flavocytochrome b558 and of several cytosolic components, including the p67^{phox}, p47^{phox}, p40^{phox} and Rac1/2 sub-units [5]. Upon phosphorylation of p47^{phox}, the different sub-units assemble to the plasma membrane forming the active enzyme. Thus, an efficient ROS production requires an optimal expression of all the NOX2 components and coordinated signals driving the enzyme assembly [5].

In this issue, Rolas and co-workers [6] add further insights into the mechanisms responsible for the deficit in neutrophils' bactericidal activity that characterizes alcoholic cirrhosis by investigating some of the mechanisms responsible for the impairment of ROS production by NOX2. They report that the deficiency in oxidative burst that characterizes the response to formylated peptides (f-MLP) of neutrophils from decompensated cirrhotic patients was associated with a severe reduction in the intracellular content of gp91^{phox}, p22^{phox} and p47^{phox}, despite their unchanged mRNA levels. By investigating the mechanisms involved they observed that the decrease of gp91^{phox} was partially due to an enhanced protein degradation due to elastase released by neutrophils themselves and more abundant in the plasma of cirrhotic patients. In addition, the lowering of NOX2 components also involved an impairment of protein synthesis consequent to the reduction of the kinase mammalian target of rapamycin (mTOR), a key regulator of cellular anabolic processes. Interestingly, treating the neutrophils from cirrhotic patients with the Toll-like receptor-7/8 (TLR-7/8) agonists CL097 and R848 restored ROS production by stimulating the synthesis of NOX2 components through a process involving mTOR activation. TLR-7/8 agonists were also active when added to whole blood, while the use of the elastase inhibitor N-(Methoxysuccinyl)-Ala-Ala-Pro-Val-chloromethylketone (NEI) had a minimal impact on NOX2 recovery. The effect of TLR agonists was rather specific, as the TLR-4 stimulation by lipopolysaccharide enhanced gp91^{phox} expression, but modestly improved ROS production. Such a specificity might likely relate to the capacity of TLR-7/8-mediated signals to potentiate NOX2 activity through the phosphorylation of p47^{phox} by protein kinase B (PKB; AKT), mTOR and p38 mitogen activated kinase (p38MAPK) [7] (Fig. 1). From these data, the authors conclude that elastase-dependent degradation and mTOR impairment contributes to the decrease gp91^{phox} and ROS production observed in patients with alcoholic cirrhosis. However, the latter process appears be more relevant to explain the defect in ROS production, as protein synthesis can compensate for the loss of gp91^{phox} due to elastase-mediated degradation. A deficit of gp91^{phox} also accounts for the lowering in ROS-mediated bacterial killing by monocytes of patients with alcoholic hepatitis [8], indicating that in alcohol abusers similar mechanisms can be responsible for affecting the functions of different phagocytic cells. Interestingly, TLR-7/8 stimulation of AKT/mTOR also contributes to restore neutrophil's bactericidal activity by promoting myeloperoxidase release [9] (Fig. 1). Nonetheless, several issues remain still open. For instance, little is known about the mechanisms that lead to the lowering of neutrophil mTOR in advanced ALD and it is still unclear how TLR-7/8 stimulation acts in reverting mTOR activity. On the other hand, the

presented data add to the consensus that mTOR inhibitors are most likely ill-advised as anti-fibrotic treatment in advanced liver fibrosis, as they will inevitably worsen the defective oxidative burst capacity of neutrophils and thus increase the risk of infections.

In conclusion, this work nicely supports previous data from the same group pointing to a key role of AKT/mTOR in the mechanisms affecting the bactericidal activity of neutrophils in alcoholic cirrhosis [7,9]. The significance of the presented data mainly derives from its possible clinical translation into using TLR-7/8 agonists to reverse alcohol-mediated impairment of neutrophil ROS production, and thus improve the host's antibacterial defence of patients with alcoholic (and non-alcoholic?) cirrhosis. Whether this could be implemented in a treatment of established infections or in their prevention remains to be determined. However, as a novel therapeutic target, TLRs clearly deserve further investigation in humans, particularly, since prophylactic antibiotic treatment in patients with cirrhosis achieves only a moderate benefit [10]. In this regard, it seems important to emphasize that TLR agonists have been shown to be effective in promoting both innate and adaptive immunity [11] and their use might exert a synergistic therapeutic effect with that of granulocyte colony-stimulating factor [12] in controlling infective complications of advanced ALD.

References

1. Albano E. Oxidative stress in alcoholic liver disease: Oxidative Stress in Applied Basic Research and Clinical Practice: Studies on Hepatic Disorders. Albano E, Parola M Eds. Humana Press, Springer International Publishing, 2015:215-239,
2. Louvet A, Mathurin P. Alcoholic liver disease: mechanisms of injury and targeted treatment. Nat Rev Gastroenterol Hepatol 2015;12:231-242.
3. Jalan R, Fernandez J, Wiest R, et al. Bacterial infections in cirrhosis: a position statement based on the EASL Special Conference 2013. J Hepatol. 2013;60:1310-1324.
4. Arroyo V, Moreau R, Jalan R et al. Acute-on-chronic liver failure: A new syndrome that will re-classify cirrhosis. J Hepatol. 2015;62 (Suppl 1):S131-S143.
5. Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. Physiol Rev 2007; 87:245-313.
6. Rolas L, Boussif A, Weiss E, et al. NADPH Oxidase depletion in neutrophils from patients with cirrhosis and restoration via Toll-like receptor 7/8 activation. Gut 2017; pii:gutjnl-2016-313443R1
7. Rolas L, Makhezer N, Hadjoudj S, et al. Inhibition of mammalian target of rapamycin aggravates the respiratory burst defect of neutrophils from decompensated patients with cirrhosis. Hepatology. 2013;57:1163-1171.
8. Vergis N, Khamri W, Beale K, et al. Defective monocyte oxidative burst predicts infection in alcoholic hepatitis and is associated with reduced expression of NADPH oxidase. Gut. 2017; 66:519-529.
9. Boussif A, Rolas L, Weiss E, et al. Impaired intracellular signaling, myeloperoxidase release and bactericidal activity of neutrophils from patients with alcoholic cirrhosis. J Hepatol. 2016;64:1041-1048.
10. Moreau R, Elkrif L, Bureau C, et al. A randomized trial of 6-month norfloxacin therapy in patients with Child-Pugh class C cirrhosis. J Hepatol 2017;66 (Suppl 1):S13
11. Vasilakos JP, Tomai MA. The use of Toll-like receptor 7/8 agonists as vaccine adjuvants. Expert Rev Vaccines. 2013;12:809-819

12. Singh V, Sharma AK, Narasimhan RL, et al. Granulocyte colony-stimulating factor in severe alcoholic hepatitis: a randomized pilot study. *Am J Gastroenterol*. 2014;109:1417-1423.

Figure caption

Figure 1: Proposed mechanisms by which TLR-7/8 stimulation improves the antibacterial activity of neutrophils from patients with alcoholic cirrhosis. Abbreviations: AKT protein kinase B; NOX2 NADPH oxidase 2; MPO myeloperoxidase; mTOR mammalian target of rapamycin; TLR-7/8 Toll-like receptor-7/8; O_2^- superoxide anion; H_2O_2 hydrogen peroxide; HClO hypochlorous acid; p38MAPK p38 mitogen activated kinase; NOX2 NADPH oxidase 2; gp91^{phox}, p22^{phox}, p67^{phox}, p47^{phox}, p40^{phox}, Rac1/2 NOX2 sub-units.




Figure 1: Proposed mechanisms by which TLR-7/8 stimulation improves the antibacterial activity of neutrophils from patients with alcoholic cirrhosis. Abbreviations: AKT protein kinase B; NOX2 NADPH oxidase 2; MPO myeloperoxidase; mTOR mammalian target of rapamycin; TLR-7/8 Toll-like receptor-7/8; O₂⁻ superoxide anion; H₂O₂ hydrogen peroxide; HClO hypochlorous acid; p38MAPK p38 mitogen activated kinase; NOX2 NADPH oxidase 2; gp91^{phox}, p22^{phox}, p67^{phox}, p47^{phox}, p40^{phox}, Rac1/2 NOX2 sub-units.

338x190mm (96 x 96 DPI)

Albano & Stickel

toll-like receptor targeting in alcoholic cirrhosis

COMMENTARY

TARGETING TOLL-LIKE RECEPTOR 7/8 IMPROVES HOST ANTI-INFECTIVE RESPONSE IN ALCOHOLIC CIRRHOSIS

Emanuele Albano¹ & Felix Stickel²

¹Department of Health Sciences and Interdisciplinary Research, Centre for Autoimmune Diseases, University of East Piedmont, Novara, Italy

²Department of Gastroenterology and Hepatology, University Hospital of Zurich, Zurich, Switzerland

Word count: 875

References: 12

Short title: Toll-like receptor targeting in alcoholic cirrhosis

Correspondence

Emanuele Albano, Dept. of Health Sciences, University of East Piedmont, Via Solaroli 17, 28100 Novara, Italy. Tel: +390321660642; Fax: +390321620421; E-mail: emanuele.albano@med.uniupo.it

The production of reactive oxygen species (ROS) by activated macrophages and neutrophil granulocytes represents an important cause of alcohol-induced oxidative stress and significantly contributes to the pathogenesis of alcoholic liver disease (ALD) [1]. Furthermore, diffuse neutrophil infiltration of the liver is a key feature of acute alcoholic hepatitis actively participating to parenchymal injury [2]. However, during ALD progression to cirrhosis, neutrophil functions such as ROS production, bacterial phagocytosis and granule exocytosis are impaired leading to an increased susceptibility to bacterial infections of cirrhotic patients [3]. Indeed, the development of bacterial peritonitis is a common complication of cirrhosis, while sepsis is a major cause of mortality of patients with decompensated alcoholic cirrhosis being also associated with multi-organ failure and immune deficiencies [4].

The key role of neutrophils in anti-bacterial defences mainly relies on the capacity of the enzyme NADPH oxidase 2 (NOX2) to generate superoxide anion (O_2^-) during a process known as oxidative burst. Superoxide anion, in fact, by conversion to hydrogen peroxide fuels myeloperoxidase-mediated production of the powerful bactericidal agents, hypochlorite and chloramine [5]. NOX2 is a multi-protein enzymatic system consisting of the transmembrane proteins gp91^{phox} and p22^{phox} forming the catalytic unit flavocytochrome b558 and of several cytosolic components, including the p67^{phox}, p47^{phox}, p40^{phox} and Rac1/2 sub-units [5]. Upon phosphorylation of p47^{phox}, the different sub-units assemble to the plasma membrane forming the active enzyme. Thus, an efficient ROS production requires an optimal expression of all the NOX2 components and coordinated signals driving the enzyme assembly [5].

In this issue, Rolas and co-workers [6] add further insights into the mechanisms responsible for the deficit in neutrophils' bactericidal activity that characterizes alcoholic cirrhosis by investigating some of the mechanisms responsible for the impairment of ROS production by NOX2. They report that the deficiency in oxidative burst that characterizes the response to formylated peptides (f-MLP) of neutrophils from decompensated cirrhotic patients was associated with a severe reduction in the intracellular content of gp91^{phox}, p22^{phox} and p47^{phox}, despite their unchanged mRNA levels. By investigating the mechanisms involved they observed that the decrease of gp91^{phox} was partially due to an enhanced protein degradation due to elastase released by neutrophils themselves and more abundant in the plasma of cirrhotic patients. In addition, the lowering of NOX2 components also involved an impairment of protein synthesis consequent to the reduction of the kinase mammalian target of rapamycin (mTOR), a key regulator of cellular anabolic processes. Interestingly, treating the neutrophils from cirrhotic patients with the Toll-like receptor-7/8 (TLR-7/8) agonists CL097 and R848 restored ROS production by stimulating the synthesis of NOX2 components through a process involving mTOR activation. TLR-7/8 agonists were also active when added to whole blood, while the use of the elastase inhibitor N-(Methoxysuccinyl)-Ala-Ala-Pro-Val-chloromethylketone (NEI) had a minimal impact on NOX2 recovery. The effect of TLR agonists was rather specific, as the TLR-4 stimulation by lipopolysaccharide enhanced gp91^{phox} expression, but modestly improved ROS production. Such a specificity might likely relate to the capacity of TLR-7/8-mediated signals to potentiate NOX2 activity through the phosphorylation of p47^{phox} by protein kinase B (PKB; AKT), mTOR and p38 mitogen activated kinase (p38MAPK) [7] (Fig. 1). From these data, the authors conclude that elastase-dependent degradation and mTOR impairment contributes to the decrease gp91^{phox} and ROS production observed in patients with alcoholic cirrhosis. However, the latter process appears be more relevant to explain the defect in ROS production, as protein synthesis can compensate for the loss of gp91^{phox} due to elastase-mediated degradation. A deficit of gp91^{phox} also accounts for the lowering in ROS-mediated bacterial killing by monocytes of patients with alcoholic hepatitis [8], indicating that in alcohol abusers similar mechanisms can be responsible for affecting the functions of different phagocytic cells. Interestingly, TLR-7/8 stimulation of AKT/mTOR also contributes to restore neutrophil's bactericidal activity by promoting myeloperoxidase release [9] (Fig. 1). Nonetheless, several issues remain still open. For instance, little is known about the mechanisms that lead to the lowering of neutrophil mTOR in advanced ALD and it is still unclear how TLR-7/8 stimulation acts in reverting mTOR activity. On the other hand, the

presented data add to the consensus that mTOR inhibitors are most likely ill-advised as anti-fibrotic treatment in advanced liver fibrosis, as they will inevitably worsen the defective oxidative burst capacity of neutrophils and thus increase the risk of infections.

In conclusion, this work nicely supports previous data from the same group pointing to a key role of AKT/mTOR in the mechanisms affecting the bactericidal activity of neutrophils in alcoholic cirrhosis [7,9]. The significance of the presented data mainly derives from its possible clinical translation into using TLR-7/8 agonists to reverse alcohol-mediated impairment of neutrophil ROS production, and thus improve the host's antibacterial defence of patients with alcoholic (and non-alcoholic?) cirrhosis. Whether this could be implemented in a treatment of established infections or in their prevention remains to be determined. However, as a novel therapeutic target, TLRs clearly deserve further investigation in humans, particularly, since prophylactic antibiotic treatment in patients with cirrhosis achieves only a moderate benefit [10]. In this regard, it seems important to emphasize that TLR agonists have been shown to be effective in promoting both innate and adaptive immunity [11] and their use might exert a synergistic therapeutic effect with that of granulocyte colony-stimulating factor [12] in controlling infective complications of advanced ALD.

References

1. Albano E. Oxidative stress in alcoholic liver disease: Oxidative Stress in Applied Basic Research and Clinical Practice: Studies on Hepatic Disorders. Albano E, Parola M Eds. Humana Press, Springer International Publishing, 2015:215-239,
2. Louvet A, Mathurin P. Alcoholic liver disease: mechanisms of injury and targeted treatment. *Nat Rev Gastroenterol Hepatol* 2015;12:231-242.
3. Jalan R, Fernandez J, Wiest R, et al. Bacterial infections in cirrhosis: a position statement based on the EASL Special Conference 2013. *J Hepatol*. 2013;60:1310-1324.
4. Arroyo V, Moreau R, Jalan R et al. Acute-on-chronic liver failure: A new syndrome that will re-classify cirrhosis. *J Hepatol*. 2015;62 (Suppl 1):S131-S143.
5. Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 2007; 87:245-313.
6. Rolas L, Boussif A, Weiss E, et al. NADPH Oxidase depletion in neutrophils from patients with cirrhosis and restoration via Toll-like receptor 7/8 activation. *Gut* 2017; pii:gutjnl-2016-313443R1
7. Rolas L, Makhezer N, Hadjoudj S, et al. Inhibition of mammalian target of rapamycin aggravates the respiratory burst defect of neutrophils from decompensated patients with cirrhosis. *Hepatology*. 2013;57:1163-1171.
8. Vergis N, Khamri W, Beale K, et al. Defective monocyte oxidative burst predicts infection in alcoholic hepatitis and is associated with reduced expression of NADPH oxidase. *Gut*. 2017; 66:519-529.
9. Boussif A, Rolas L, Weiss E, et al. Impaired intracellular signaling, myeloperoxidase release and bactericidal activity of neutrophils from patients with alcoholic cirrhosis. *J Hepatol*. 2016;64:1041-1048.
10. Moreau R, Elkrif L, Bureau C, et al. A randomized trial of 6-month norfloxacin therapy in patients with Child-Pugh class C cirrhosis. *J Hepatol* 2017;66 (Suppl 1):S13
11. Vasilakos JP, Tomai MA. The use of Toll-like receptor 7/8 agonists as vaccine adjuvants. *Expert Rev Vaccines*. 2013;12:809-819

12. Singh V, Sharma AK, Narasimhan RL, et al. Granulocyte colony-stimulating factor in severe alcoholic hepatitis: a randomized pilot study. Am J Gastroenterol. 2014;109:1417-1423.

Figure caption

Figure 1: Proposed mechanisms by which TLR-7/8 stimulation improves the antibacterial activity of neutrophils from patients with alcoholic cirrhosis. Abbreviations: AKT protein kinase B; NOX2 NADPH oxidase 2; MPO myeloperoxidase; mTOR mammalian target of rapamycin; TLR-7/8 Toll-like receptor-7/8; O₂⁻ superoxide anion; H₂O₂ hydrogen peroxide; HClO hypochlorous acid; p38MAPK p38 mitogen activated kinase; NOX2 NADPH oxidase 2; gp91^{phox}, p22^{phox}, p67^{phox}, p47^{phox}, p40^{phox}, Rac1/2 NOX2 sub-units.